

GLOBAL SOLVABILITY OF 3D INHOMOGENEOUS NAVIER-STOKES EQUATIONS WITH DENSITY-DEPENDENT VISCOSITY

XIANGDI HUANG AND YUN WANG

ABSTRACT. In this paper, we consider the three-dimensional inhomogeneous Navier-Stokes equations with density-dependent viscosity in presence of vacuum over bounded domains. Global-in-time unique strong solution is proved to exist when $\|\nabla u_0\|_{L^2}$ is suitably small with arbitrary large initial density. This generalizes all the previous results even for the constant viscosity.

Keywords: density-dependent viscosity, inhomogeneous Navier-Stokes equations, strong solution, vacuum.

AMS: 35Q35, 35B65, 76N10

1. INTRODUCTION

The Navier-Stokes equations are usually used to describe the motion of fluids. In particular, for the study of multiphase fluids without surface tension, the following density-dependent Navier-Stokes equations acts as a model on some bounded domain $\Omega \subset R^N (N = 2, 3)$,

$$(1.1) \quad \begin{cases} \rho_t + \operatorname{div}(\rho u) = 0, & \text{in } \Omega \times (0, T], \\ (\rho u)_t + \operatorname{div}(\rho u \otimes u) - \operatorname{div}(2\mu(\rho)d) + \nabla P = 0, & \text{in } \Omega \times (0, T], \\ \operatorname{div} u = 0, & \text{in } \Omega \times [0, T], \\ u = 0, & \text{on } \partial\Omega \times [0, T], \\ \rho|_{t=0} = \rho_0, \quad u|_{t=0} = u_0, & \text{in } \Omega. \end{cases}$$

Here ρ , u , and P denote the density, velocity and pressure of the fluid, respectively.

$$d = \frac{1}{2} [\nabla u + (\nabla u)^T]$$

is the deformation tensor.

$\mu = \mu(\rho)$ states the viscosity and is a function of ρ , which is assumed to satisfy

$$(1.2) \quad \mu \in C^1[0, \infty), \quad \text{and} \quad \mu \geq \underline{\mu} > 0 \quad \text{on } [0, \infty) \quad \text{for some positive constant } \underline{\mu}.$$

In this paper, we study the initial boundary value problem to the system (1.1)-(1.2).

The mathematical study for nonhomogeneous incompressible flow was initiated by the Russian school. They studied the case that $\mu(\rho)$ is a constant and the initial density ρ_0 is bounded away from 0. In the absence of vacuum, global existence of weak solutions as well as local strong solution was established by Kazhikov [4, 21]. The uniqueness of local strong solutions was first established by Ladyzhenskaya-Solonnikov [22] for the initial boundary value problem, see also [25]. Furthermore, unique local strong solution is proved to be global in

2D [26]. In recent years, Danchin initiated the studies for solutions in critical spaces. He [9, 10] derived the global well-posedness for small initial velocity in critical spaces, where density is close to a constant. For some subsequent works, refer to [1, 24] and references therein. We remark that in the very interesting papers [11, 12], Danchin-Mucha studied the case for which density is piecewise constant, see also some generalizations in 2D [18].

When initial vacuum is taken into account and $\mu(\rho)$ is still a constant, Simon [26] proved the global existence of weak solutions. Later, Choe-Kim [7] proposed a compatibility condition as (1.4) below to establish local existence of strong solution. Global strong solution allowing vacuum in 2D was recently derived by the authors [20]. Meanwhile, some global solutions in 3D with small critical norms have been constructed, refer to the results in [2, 8] and references therein.

Finally, we come to the most general case: viscosity $\mu(\rho)$ depends on density ρ . Most results were concentrated on 2D case. Global weak solutions were derived by the revolutionary work [14, 23] of DiPerna and Lions. Later, Desjardins [13] proved the global weak solution with more regularity for the two-dimensional case provided that the viscosity function $\mu(\rho)$ is a small perturbation of a positive constant in L^∞ - norm. Very recently, Abidi-Zhang [3] generalized this 2D result to strong solutions. Regarding the 3D case, Cho-Kim [6] constructed a unique local strong solution by imposing some initial compatibility condition. Their result is stated as follows:

Theorem 1.1. *Assume that the initial data (ρ_0, u_0) satisfies the regularity condition*

$$(1.3) \quad 0 \leq \rho_0 \in W^{1,q}, \quad 3 < q < \infty, \quad u_0 \in H_{0,\sigma}^1 \cap H^2,$$

and the compatibility condition

$$(1.4) \quad -\operatorname{div} (\mu(\rho_0) [\nabla u_0 + (\nabla u_0)^T]) + \nabla P_0 = \rho_0^{\frac{1}{2}} g,$$

for some $(P_0, g) \in H^1 \times L^2$. Then there exists a small time T and a unique strong solution (ρ, u, P) to the initial boundary value problem (1.1) such that

$$\begin{aligned} \rho &\in C([0, T]; W^{1,q}), & \nabla u, P &\in C([0, T]; H^1) \cap L^2(0, T; W^{1,r}), \\ \rho_t &\in C([0, T]; L^q), & \sqrt{\rho} u_t &\in L^\infty(0, T; L^2), \quad u_t \in L^2(0, T; H_0^1), \end{aligned}$$

for any r with $1 \leq r < q$. Furthermore, if T^* is the maximal existence time of the local strong solution (ρ, u) , then either $T^* = \infty$ or

$$(1.5) \quad \sup_{0 \leq t < T^*} (\|\nabla \rho(t)\|_{L^q} + \|\nabla u(t)\|_{L^2}) = \infty.$$

Motivated by the global existence result [8] for the special case that μ is a constant, we aim to establish global well-posedness result for variable coefficient case. However, due to the strong coupling between viscosity coefficient and density, it's more complicated and involved with variable coefficient $\mu(\rho)$ and requires more delicate analysis.

Our main result proves the existence of global strong solution, provided $\|\nabla u_0\|_{L^2}$ is suitably small. To conclude, we arrive at

Theorem 1.2. *Assume that the initial data (ρ_0, u_0) satisfies (1.3)-(1.4), and $0 \leq \rho_0 \leq \bar{\rho}$. Then there exists some small positive constant ϵ_0 , depending on Ω , q , $\bar{\rho}$, $\bar{\mu} = \sup_{[0, \bar{\rho}]} \mu(\rho)$, $\underline{\mu}$, $\|\nabla \mu(\rho_0)\|_{L^q} (= M)$, such that if*

$$(1.6) \quad \|\nabla u_0\|_{L^2} \leq \epsilon_0,$$

then the initial boundary value problem (1.1) admits a unique global strong solution (ρ, u) , with

$$(1.7) \quad \begin{aligned} \rho &\in C([0, \infty); W^{1,q}), & \nabla u, P &\in C([0, \infty); H^1) \cap L_{loc}^2(0, \infty; W^{1,r}), \\ \rho_t &\in C([0, \infty); L^q), & \sqrt{\rho} u_t &\in L_{loc}^\infty(0, \infty; L^2), \quad u_t \in L_{loc}^2(0, \infty; H_0^1), \end{aligned}$$

for any r , $1 \leq r < q$.

The main idea is to combine techniques developed by the authors in [8, 19] and time weighted energy estimates successfully applied to compressible Navier-Stokes equation by Hoff [17].

Let's briefly sketch the proof. First we assume that $\|\nabla \mu(\rho)\|_{L^q}$ is less than $4M$ and $\|\nabla u\|_{L^2}^2$ is less than $4\|\nabla u_0\|_{L^2}^2$ on $[0, T]$, then we prove that in fact $\|\nabla \mu(\rho)\|_{L^q}$ is less than $2M$ and $\|\nabla u\|_{L^2}^2$ is less than $2\|\nabla u_0\|_{L^2}^2$ on $[0, T]$, under the assumption $\|\nabla u_0\|_{L^2} \leq \epsilon_0 \leq \frac{1}{2}$. On the other hand, the control of $\|\nabla \mu(\rho)\|_{L^q}$ and $\|\nabla u\|_{L^2}$ lead to uniform estimates for other quantities, which guarantee the extension of local strong solutions. All the above procedures depends on a time independent bound of $\|\nabla u\|_{L^1 L^\infty}$. Thanks to bounded domain, u indeed has exponentially decay rather than insufficient polynomial decay for the whole space, that's the main reason why we can only treat system (1.1) in bounded domain.

Remark 1.1. *Theorem 1.2 also hold for 2D case. Since the proof is quite similar, we omit it for simplicity.*

The rest of the paper is organized as follows: Section 2 consists of some notations, definitions, and basic lemmas. Section 3 is devoted to the proof of Theorem 1.2.

2. PRELIMINARIES

Ω is a bounded smooth domain in $\mathbb{R}^3$. Denote

$$\int f dx = \int_{\Omega} f dx.$$

For $1 \leq r \leq \infty$ and $k \in \mathbb{N}$, the Sobolev spaces are defined in a standard way,

$$L^r = L^r(\Omega), \quad W^{k,r} = \{f \in L^r : \nabla^k f \in L^r\},$$

$$H^k = W^{k,2}, \quad C_{0,\sigma}^\infty = \{f \in C_0^\infty : \operatorname{div} f = 0\}.$$

$$H_0^1 = \overline{C_0^\infty}, \quad H_{0,\sigma}^1 = \overline{C_{0,\sigma}^\infty}, \quad \text{closure in the norm of } H^1.$$

High-order a priori estimates rely on the following regularity results for density-dependent Stokes equations.

Lemma 2.1. *Assume that $\rho \in W^{1,q}$, $3 < q < \infty$, and $0 \leq \rho \leq \bar{\rho}$. Let $(u, P) \in H_{0,\sigma}^1 \times L^2$ be the unique weak solution to the boundary value problem*

$$(2.8) \quad -\operatorname{div} (2\mu(\rho)d) + \nabla P = F, \quad \operatorname{div} u = 0 \quad \text{in } \Omega, \quad \text{and} \quad \int \frac{P}{\mu(\rho)} dx = 0,$$

where $d = \frac{1}{2} [\nabla u + (\nabla u)^T]$ and

$$\mu \in C^1[0, \infty), \quad \underline{\mu} \leq \mu(\rho) \leq \bar{\mu} \quad \text{on } [0, \bar{\rho}].$$

Then we have the following regularity results:

(1) If $F \in L^2$, then $(u, P) \in H^2 \times H^1$ and

$$(2.9) \quad \|u\|_{H^2} + \|P/\mu(\rho)\|_{H^1} \leq C \left(\frac{1}{\underline{\mu}} + \frac{\bar{\mu}}{\underline{\mu}^{\frac{1}{\theta_2}+2}} \|\nabla \mu\|_{L^q}^{\frac{1}{\theta_2}} \right) \|F\|_{L^2},$$

where θ_2 satisfies

$$\frac{1}{2} - \frac{1}{q} = \frac{\theta_2}{3} + \frac{1}{6}, \quad \text{i. e.,} \quad \theta_2 = \frac{q}{q-3}.$$

(2) If $F \in L^r$ for some $r \in (2, q)$, then $(u, P) \in W^{2,r} \times W^{1,r}$ and

$$(2.10) \quad \|u\|_{W^{2,r}} + \|P/\mu(\rho)\|_{W^{1,r}} \leq C \left(\frac{1}{\underline{\mu}} + \frac{\bar{\mu}}{\underline{\mu}^{\frac{1}{\theta_r}+2}} \|\nabla \mu\|_{L^q}^{\frac{1}{\theta_r}} \right) \|F\|_{L^r},$$

where

$$\frac{1}{\theta_r} = \frac{\frac{5}{6} - \frac{1}{r}}{\frac{1}{3} - \frac{1}{q}}.$$

Here the constant C in (2.9) and (2.10) depends on Ω , q , r .

The proof of Lemma 2.1 has been given in [6], although the lemma is slightly different from the version in [6]. We sketch it here for completeness.

Proof. For the existence and uniqueness of the solution, please refer to Giaquinta-Modica [16]. We give the a priori estimates here. Assume that $F \in L^2$. Multiply the first equation of (2.8) by u and integrate over Ω , then by Poincaré's inequality,

$$\int 2\mu(\rho)|d|^2 dx = \int F \cdot u dx \leq \|F\|_{L^2} \cdot \|u\|_{L^2} \leq C \|F\|_{L^2} \cdot \|\nabla u\|_{L^2}.$$

Note that

$$2 \int |d|^2 dx = \int |\nabla u|^2 dx,$$

hence

$$(2.11) \quad \|\nabla u\|_{L^2} \leq C \underline{\mu}^{-1} \|F\|_{L^2}.$$

Since $\int \frac{P}{\mu(\rho)} dx = 0$, according to Bovosgii's theory, there exists a function $v \in H_0^1$, such that

$$\operatorname{div} v = \frac{P}{\mu(\rho)},$$

and

$$\|\nabla v\|_{L^2} \leq C \left\| \frac{P}{\mu(\rho)} \right\|_{L^2}.$$

Multiply the first equation of (2.8) by $-v$, and integrate over Ω , then

$$\begin{aligned} \int \frac{P^2}{\mu(\rho)} dx &= - \int F \cdot v dx + 2 \int \mu(\rho) d : \nabla v dx \\ &\leq \|F\|_{L^2} \cdot \|v\|_{L^2} + C \bar{\mu} \cdot \|\nabla u\|_{L^2} \cdot \|\nabla v\|_{L^2} \\ &\leq C \|F\|_{L^2} \cdot \|\nabla v\|_{L^2} + C \frac{\bar{\mu}}{\underline{\mu}} \cdot \|F\|_{L^2} \cdot \|\nabla v\|_{L^2} \\ &\leq C \frac{\bar{\mu}}{\underline{\mu}} \cdot \|F\|_{L^2} \cdot \left\| \frac{P}{\mu(\rho)} \right\|_{L^2}. \end{aligned}$$

On the other hand side,

$$\int \frac{P^2}{\mu(\rho)} dx \geq \underline{\mu} \int \frac{P^2}{\mu(\rho)^2} dx.$$

Hence,

$$(2.12) \quad \left\| \frac{P}{\mu(\rho)} \right\|_{L^2} \leq C \frac{\bar{\mu}}{\underline{\mu}^2} \cdot \|F\|_{L^2}.$$

The first equation of (2.8) can be re-written as

$$-\Delta u + \nabla \left(\frac{P}{\mu(\rho)} \right) = \frac{F}{\mu(\rho)} + \frac{2d \cdot \nabla \mu(\rho)}{\mu(\rho)} - \frac{P \nabla \mu(\rho)}{\mu(\rho)^2}.$$

By virtue of the classical theory for Stokes equations and Gagliardo-Nirenberg inequality, we have

$$\begin{aligned} &\|u\|_{H^2} + \left\| \nabla \left(\frac{P}{\mu(\rho)} \right) \right\|_{L^2} \\ &\leq C \left(\left\| \frac{F}{\mu(\rho)} \right\|_{L^2} + \left\| \frac{d \cdot \nabla \mu(\rho)}{\mu(\rho)} \right\|_{L^2} + \left\| \frac{P \nabla \mu(\rho)}{\mu(\rho)^2} \right\|_{L^2} \right) \\ &\leq C \left(\underline{\mu}^{-1} \|F\|_{L^2} + \underline{\mu}^{-1} \|\nabla \mu\|_{L^q} \cdot \|\nabla u\|_{L^{\frac{2q}{q-2}}} + \underline{\mu}^{-1} \|\nabla \mu\|_{L^q} \cdot \left\| \frac{P}{\mu} \right\|_{L^{\frac{2q}{q-2}}} \right) \\ &\leq C \left[\underline{\mu}^{-1} \|F\|_{L^2} + \underline{\mu}^{-1} \|\nabla \mu\|_{L^q} \cdot \|\nabla u\|_{L^2}^{\theta_2} \cdot \|\nabla u\|_{H^1}^{1-\theta_2} \right. \\ &\quad \left. + \underline{\mu}^{-1} \|\nabla \mu\|_{L^q} \cdot \left\| \frac{P}{\mu} \right\|_{L^2}^{\theta_2} \cdot \left\| \nabla \left(\frac{P}{\mu} \right) \right\|_{L^2}^{1-\theta_2} \right]. \end{aligned}$$

By Young's inequality,

$$\begin{aligned}
& \|u\|_{H^2} + \left\| \nabla \left(\frac{P}{\mu(\rho)} \right) \right\|_{L^2} \\
& \leq C \underline{\mu}^{-1} \|F\|_{L^2} + C \underline{\mu}^{-\frac{1}{\theta_2}} \|\nabla \mu\|_{L^q}^{\frac{1}{\theta_2}} \cdot \left(\|\nabla u\|_{L^2} + \left\| \frac{P}{\mu} \right\|_{L^2} \right) \\
(2.13) \quad & \leq C \underline{\mu}^{-1} \|F\|_{L^2} + C \underline{\mu}^{-\frac{1}{\theta_2}-1} \cdot \left(1 + \frac{\bar{\mu}}{\underline{\mu}} \right) \cdot \|\nabla \mu\|_{L^q}^{\frac{1}{\theta_2}} \cdot \|F\|_{L^2} \\
& \leq C \left(\frac{1}{\underline{\mu}} + \frac{\bar{\mu}}{\underline{\mu}^{\frac{1}{\theta_2}+2}} \|\nabla \mu\|_{L^q}^{\frac{1}{\theta_2}} \right) \|F\|_{L^2},
\end{aligned}$$

where θ_2 satisfies

$$\frac{1}{2} - \frac{1}{q} = \frac{\theta_2}{3} + \frac{1}{6}, \quad \text{or} \quad \theta_2 = \frac{q}{q-3}.$$

Similarly,

$$(2.14) \quad \|u\|_{W^{2,r}} + \|\nabla (P/\mu(\rho))\|_{L^r} \leq C \left(\frac{1}{\underline{\mu}} + \frac{\bar{\mu}}{\underline{\mu}^{\frac{1}{\theta_r}+2}} \|\nabla \mu\|_{L^q}^{\frac{1}{\theta_r}} \right) \|F\|_{L^r},$$

where

$$\frac{1}{\theta_r} = \frac{\frac{5}{6} - \frac{1}{r}}{\frac{1}{3} - \frac{1}{q}}.$$

□

3. PROOF OF THEOREM 1.2

The proof of Theorem 1.2 is composed of two parts. The first part contains a priori time-weighted estimates of different levels. Upon these estimates, the second part uses a contradiction induction process to extend the local strong solution. The two parts are presented in Subsections 3.1 and 3.2, respectively.

3.1. A Priori Estimates. In this subsection, we establish some a priori time-weighted estimates. The initial velocity belongs to H^1 , but some uniform estimates of higher order and independent of time are required. To achieve that, we take some power of time as a weight. The idea is based on the parabolic property of the system. In this subsection, the constant C will denote some positive constant which maybe dependent on Ω , q , but is independent of ρ_0 or u_0 .

First, as the density satisfies the transport equation (1.1)₁ and making use of (1.1)₃, one has the following lemma.

Lemma 3.1. *Suppose (ρ, u, P) is the unique local strong solution to (1.1) on $[0, T]$, with the initial data (ρ_0, u_0) , it holds that*

$$0 \leq \rho(x, t) \leq \bar{\rho}, \quad \text{for every } (x, t) \in \Omega \times [0, T].$$

Next, the basic energy inequality of the system (1.1) reads

Theorem 3.2. *Suppose (ρ, u, P) is the unique local strong solution to (1.1) on $[0, T]$, with the initial data (ρ_0, u_0) , it holds that*

$$(3.15) \quad \int \rho |u(t)|^2 dx + \int_0^t \int \mu(\rho) |d|^2 dx ds \leq C \cdot \bar{\rho} \cdot \|u_0\|_{L^2}^2, \quad \text{for every } t \in [0, T],$$

or in other words,

$$(3.16) \quad \int \rho |u(t)|^2 dx + \underline{\mu} \int_0^t \int |\nabla u|^2 dx ds \leq C \cdot \bar{\rho} \cdot \|u_0\|_{L^2}^2, \quad \text{for every } t \in [0, T].$$

Proof. The proof is standard. Multiplying the momentum equation by u and integrating over Ω yield that

$$\frac{1}{2} \frac{d}{dt} \int \rho |u|^2 dx + 2 \int \mu(\rho) |d|^2 dx = 0.$$

Then (3.16) is true owing to the fact $2 \int |d|^2 dx = \int |\nabla u|^2 dx$ and $\mu(\rho) \geq \underline{\mu}$. $\square$

Denote

$$M = \|\nabla \mu(\rho_0)\|_{L^q},$$

$$M_2 = \frac{1}{\underline{\mu}} + \frac{\bar{\mu}}{\underline{\mu}} \cdot \frac{1}{\underline{\mu}^{1/\theta_2+1}} \cdot (4M)^{\frac{1}{\theta_2}},$$

and

$$M_r = \frac{1}{\underline{\mu}} + \frac{\bar{\mu}}{\underline{\mu}} \cdot \frac{1}{\underline{\mu}^{1/\theta_r+1}} \cdot (4M)^{\frac{1}{\theta_r}}.$$

Theorem 3.3. *Suppose (ρ, u, P) is the unique local strong solution to (1.1) on $[0, T]$, with the initial data (ρ_0, u_0) , and satisfies*

$$(3.17) \quad \sup_{t \in [0, T]} \|\nabla \mu(\rho(t))\|_{L^q} \leq 4M,$$

and

$$(3.18) \quad \sup_{t \in [0, T]} \|\nabla u(t)\|_{L^2}^2 \leq 4 \|\nabla u_0\|_{L^2}^2 \leq 1.$$

There exists a positive number C_1 , depending on Ω, q such that if

$$(3.19) \quad C_1 \underline{\mu}^{-2} (M_2^2 + M^4 M_2^6) \bar{\rho}^4 \cdot \|\nabla u_0\|_{L^2}^2 \leq \ln 2,$$

then

$$(3.20) \quad \frac{1}{\underline{\mu}} \int_0^T \|\sqrt{\rho} u_t\|_{L^2}^2 dt + \sup_{t \in [0, T]} \|\nabla u(t)\|_{L^2}^2 \leq 2 \|\nabla u_0\|_{L^2}^2.$$

Before the proof of Theorem 3.3, let us introduce an auxiliary lemma, which is a result of the $W^{2,2}$ -estimates in Lemma 2.1.

Lemma 3.4. *Suppose (ρ, u, P) is the unique local strong solution to (1.1) on $[0, T]$ and satisfies*

$$\sup_{t \in [0, T]} \|\nabla \mu(\rho(t))\|_{L^q} \leq 4M.$$

Then it holds that

$$\|\nabla u\|_{H^1} \leq C M_2 \|\rho u_t\|_{L^2} + C M_2^2 \cdot \bar{\rho}^2 \cdot \|\nabla u\|_{L^2}^3.$$

Proof. The momentum equation can be rewritten as follows,

$$(3.21) \quad -2\operatorname{div}(\mu(\rho)d) + \nabla P = -\rho u_t - (\rho u \cdot \nabla)u.$$

It follows from Lemma 2.1 and Gagliardo-Nirenberg inequality that

$$\begin{aligned} \|\nabla u\|_{H^1} &\leq CM_2(\|\rho u_t\|_{L^2} + \|\rho u \cdot \nabla u\|_{L^2}) \\ &\leq CM_2\|\rho u_t\|_{L^2} + CM_2 \cdot \bar{\rho} \cdot \|u\|_{L^6} \cdot \|\nabla u\|_{L^3} \\ &\leq CM_2\|\rho u_t\|_{L^2} + CM_2 \cdot \bar{\rho} \cdot \|\nabla u\|_{L^2}^{\frac{3}{2}} \cdot \|\nabla u\|_{H^1}^{\frac{1}{2}}. \end{aligned}$$

By Young's inequality,

$$\|\nabla u\|_{H^1} \leq CM_2\|\rho u_t\|_{L^2} + CM_2^2 \cdot \bar{\rho}^2 \cdot \|\nabla u\|_{L^2}^3.$$

□

Proof of Theorem 3.3. Multiply the momentum equation by u_t and integrate over Ω , then

$$\begin{aligned} &\int \rho |u_t|^2 dx + \frac{d}{dt} \int \mu(\rho) |d|^2 dx \\ &\leq \left| \int \rho u \cdot \nabla u \cdot u_t dx \right| + C \int |\nabla \mu(\rho)| \cdot |u| \cdot |\nabla u|^2 dx. \end{aligned}$$

Here we used the renormalized mass equation for $\mu(\rho)$,

$$\partial_t[\mu(\rho)] + u \cdot \nabla \mu(\rho) = 0,$$

which is derived due to the fact $\operatorname{div} u = 0$.

Applying Gagliardo-Nirenberg inequality and Lemma 3.4,

$$\begin{aligned} &\left| \int \rho u \cdot \nabla u \cdot u_t dx \right| \\ &\leq \frac{1}{8} \|\sqrt{\rho} u_t\|_{L^2}^2 + C \|\sqrt{\rho} u\|_{L^6}^2 \cdot \|\nabla u\|_{L^3}^2 \\ &\leq \frac{1}{8} \|\sqrt{\rho} u_t\|_{L^2}^2 + C \bar{\rho} \cdot \|\nabla u\|_{L^2}^3 \cdot \|\nabla u\|_{H^1} \\ &\leq \frac{1}{8} \|\sqrt{\rho} u_t\|_{L^2}^2 + C \bar{\rho} \cdot \|\nabla u\|_{L^2}^3 \cdot [CM_2\|\rho u_t\|_{L^2} + CM_2^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^3], \end{aligned}$$

and similarly,

$$\begin{aligned} &C \int |\nabla \mu(\rho)| \cdot |u| \cdot |\nabla u|^2 dx \\ &\leq C \|\nabla \mu(\rho)\|_{L^3} \cdot \|u\|_{L^6} \cdot \|\nabla u\|_{L^4}^2 \\ &\leq C \|\nabla \mu(\rho)\|_{L^3} \cdot \|\nabla u\|_{L^2}^{\frac{3}{2}} \cdot \|\nabla u\|_{H^1}^{\frac{3}{2}} \\ &\leq CM \|\nabla u\|_{L^2}^{\frac{3}{2}} \cdot [CM_2\|\rho u_t\|_{L^2} + CM_2^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^3]^{\frac{3}{2}}. \end{aligned}$$

Hence, by Young's inequality,

$$\begin{aligned}
& \int \rho |u_t|^2 dx + \frac{d}{dt} \int \mu(\rho) |d|^2 dx \\
& \leq \frac{1}{8} \|\sqrt{\rho} u_t\|_{L^2}^2 + \frac{1}{8} \|\sqrt{\rho} u_t\|_{L^2}^2 + C M_2^2 \bar{\rho}^3 \|\nabla u\|_{L^2}^6 + \frac{1}{8} \|\sqrt{\rho} u_t\|_{L^2}^2 \\
& \quad + C \left(M M_2^{\frac{3}{2}} \cdot \bar{\rho}^{\frac{3}{4}} \|\nabla u\|_{L^2}^{\frac{3}{2}} \right)^4 + C M M_2^3 \cdot \bar{\rho}^3 \cdot \|\nabla u\|_{L^2}^6 \\
& \leq \frac{3}{8} \|\sqrt{\rho} u_t\|_{L^2}^2 + C (M_2^2 + M^4 M_2^6 + M M_2^3) \cdot \bar{\rho}^3 \cdot \|\nabla u\|_{L^2}^6.
\end{aligned}$$

So we have

$$(3.22) \quad \int \rho |u_t|^2 dx + \frac{d}{dt} \int \mu(\rho) |d|^2 dx \leq C (M_2^2 + M^4 M_2^6) \cdot \bar{\rho}^3 \cdot \|\nabla u\|_{L^2}^6.$$

Integrate with respect to time on $[0, t]$,

$$\frac{1}{\underline{\mu}} \int_0^t \int \rho |u_t|^2 dx ds + \sup_{s \in [0, t]} \int |\nabla u|^2 dx \leq C \underline{\mu}^{-1} (M_2^2 + M^4 M_2^6) \cdot \bar{\rho}^3 \cdot \int_0^t \|\nabla u\|_{L^2}^6 ds.$$

Applying Gronwall's inequality,

$$\begin{aligned}
& \frac{1}{\underline{\mu}} \int_0^T \int \rho |u_t|^2 dx dt + \sup_{t \in [0, T]} \int |\nabla u|^2 dx \\
& \leq \|\nabla u_0\|_{L^2}^2 \cdot \exp \left\{ C \underline{\mu}^{-1} (M_2^2 + M^4 M_2^6) \cdot \bar{\rho}^3 \cdot \int_0^T \|\nabla u\|_{L^2}^4 dt \right\}.
\end{aligned}$$

According to Theorem 3.2 and the assumption (3.18),

$$\begin{aligned}
(3.23) \quad & \int_0^T \|\nabla u\|_{L^2}^4 dt \leq \sup_{t \in [0, T]} \|\nabla u\|_{L^2}^2 \cdot \int_0^T \|\nabla u\|_{L^2}^2 ds \\
& \leq C \underline{\mu}^{-1} \cdot \bar{\rho} \cdot \|u_0\|_{L^2}^2 \\
& \leq C \underline{\mu}^{-1} \cdot \bar{\rho} \cdot \|\nabla u_0\|_{L^2}^2.
\end{aligned}$$

Hence, we arrive at

$$\begin{aligned}
(3.24) \quad & \frac{1}{\underline{\mu}} \int_0^T \int \rho |u_t|^2 dx dt + \sup_{t \in [0, T]} \int |\nabla u|^2 dx \\
& \leq \|\nabla u_0\|_{L^2}^2 \exp \{ C_1 \underline{\mu}^{-2} (M_2^2 + M^4 M_2^6) \cdot \bar{\rho}^4 \cdot \|\nabla u_0\|_{L^2}^2 \}.
\end{aligned}$$

Now it is clear that (3.20) holds, provided (3.19) holds.

□

As a byproduct of the estimates in the proof, we have the following result.

Theorem 3.5. *Suppose (ρ, u, P) is the unique local strong solution to (1.1) on $[0, T]$, with the initial data (ρ_0, u_0) , and satisfies the assumptions (3.17)-(3.19) as in Theorem 3.3. Then*

$$(3.25) \quad \frac{1}{\underline{\mu}} \int_0^T t \|\sqrt{\rho} u_t\|_{L^2}^2 dt + \sup_{t \in [0, T]} t \|\nabla u\|_{L^2}^2 \leq \frac{C \cdot \bar{\rho}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2.$$

Proof. Multiplying (3.22) by t , as shown in the last proof, one has

$$(3.26) \quad \begin{aligned} & \frac{1}{\underline{\mu}} \int_0^T t \|\sqrt{\rho} u_t\|_{L^2}^2 dt + \sup_{t \in [0, T]} t \|\nabla u\|_{L^2}^2 \\ & \leq \frac{1}{\underline{\mu}} \int_0^T \int \mu(\rho) |d|^2 dx dt \cdot \exp\{C_1 \underline{\mu}^{-2} (M_2^2 + M^4 M_2^6) \cdot \bar{\rho}^4 \cdot \|\nabla u_0\|_{L^2}^2\}. \end{aligned}$$

According to Theorem 3.2,

$$(3.27) \quad \int_0^T \int \mu(\rho) |d|^2 dx dt \leq C \cdot \bar{\rho} \cdot \|u_0\|_{L^2}^2 \leq C \cdot \bar{\rho} \cdot \|\nabla u_0\|_{L^2}^2.$$

Hence,

$$\begin{aligned} & \frac{1}{\underline{\mu}} \int_0^T t \|\sqrt{\rho} u_t\|_{L^2}^2 dt + \sup_{t \in [0, T]} t \|\nabla u\|_{L^2}^2 \\ & \leq \frac{C \bar{\rho}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2 \cdot \exp\{C_1 \underline{\mu}^{-2} (M_2^2 + M^2 M_2^6) \cdot \bar{\rho}^4 \cdot \|\nabla u_0\|_{L^2}^2\} \\ & \leq \frac{C \cdot \bar{\rho}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2. \end{aligned}$$

□

Theorem 3.6. *Suppose (ρ, u, P) is the unique local strong solution to (1.1) on $[0, T]$, with the initial data (ρ_0, u_0) , and satisfies the assumptions (3.17)-(3.19). Then*

$$(3.28) \quad \begin{aligned} & \sup_{t \in [0, T]} t \int \rho |u_t|^2 dx + \underline{\mu} \int_0^T t \|\nabla u_t\|_{L^2}^2 \\ & \leq C \|\nabla u_0\|_{L^2}^2 \Theta_1 \cdot \exp\{C \Theta_2\} \end{aligned}$$

and

$$(3.29) \quad \begin{aligned} & \sup_{t^2 \in [0, T]} t^2 \int \rho |u_t|^2 dx + \underline{\mu} \int_0^T t^2 \|\nabla u_t\|_{L^2}^2 \\ & \leq C \frac{\bar{\rho}}{\underline{\mu}} \|\nabla u_0\|_{L^2}^2 \Theta_1 \cdot \exp\{C \Theta_2\}. \end{aligned}$$

where

$$(3.30) \quad \Theta_1 = \frac{M_2^4 \bar{\rho}^8}{\underline{\mu}^3} + \frac{M^2 M_2^8 \bar{\rho}^{10}}{\underline{\mu}^3} + \underline{\mu}, \quad \Theta_2 = \frac{\bar{\rho}^4}{\underline{\mu}^4} + \frac{M_2^2 \bar{\rho}^4}{\underline{\mu}^2} + M^2 M_2^4 \bar{\rho}^2.$$

Proof. Take t -derivative of the momentum equation,

$$(3.31) \quad \rho u_{tt} + (\rho u) \cdot \nabla u_t - \operatorname{div} (2\mu(\rho) d_t) + \nabla P_t = -\rho_t u_t - (\rho u)_t \cdot \nabla u + \operatorname{div} (2\mu(\rho)_t d).$$

Multiplying (3.31) by tu_t and integrating over Ω , we get after integration by parts that

$$(3.32) \quad \begin{aligned} & \frac{t}{2} \frac{d}{dt} \int \rho |u_t|^2 dx + 2t \int \mu(\rho) |d_t|^2 dx \\ & = -t \int \rho_t |u_t|^2 dx - t \int (\rho u)_t \cdot \nabla u \cdot u_t dx - t \int 2\mu(\rho)_t d \cdot \nabla u_t dx. \end{aligned}$$

Let us estimate the terms on the righthand of (3.32). First, utilizing the mass equation and Poincaré's inequality, one has

$$\begin{aligned}
(3.33) \quad & -t \int \rho_t |u_t|^2 dx \\
& = -2t \int \rho u \cdot \nabla u_t \cdot u_t dx \\
& \leq C \bar{\rho}^{\frac{1}{2}} \cdot t \cdot \|\sqrt{\rho} u_t\|_{L^3} \cdot \|\nabla u_t\|_{L^2} \cdot \|u\|_{L^6} \\
& \leq C \bar{\rho}^{\frac{1}{2}} \cdot t \cdot \|\sqrt{\rho} u_t\|_{L^2}^{\frac{1}{2}} \cdot \|\sqrt{\rho} u_t\|_{L^6}^{\frac{1}{2}} \cdot \|\nabla u_t\|_{L^2} \cdot \|\nabla u\|_{L^2} \\
& \leq C \bar{\rho}^{\frac{3}{4}} \cdot t \cdot \|\sqrt{\rho} u_t\|_{L^2}^{\frac{1}{2}} \cdot \|\nabla u_t\|_{L^2}^{\frac{3}{2}} \cdot \|\nabla u\|_{L^2} \\
& \leq \frac{1}{8} \underline{\mu} \cdot t \|\nabla u_t\|_{L^2}^2 + C \underline{\mu}^{-3} \cdot \bar{\rho}^3 \cdot t \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4.
\end{aligned}$$

Second, utilizing the renormalized mass equation for $\mu(\rho)$,

$$\begin{aligned}
(3.34) \quad & -2t \int \mu(\rho)_t \cdot d \cdot \nabla u_t dx \\
& \leq Ct \int |u| \cdot |\nabla \mu(\rho)| \cdot |d| \cdot |\nabla u_t| dx \\
& \leq Ct \cdot \|\nabla \mu(\rho)\|_{L^3} \cdot \|\nabla u_t\|_{L^2} \cdot \|d\|_{L^6} \cdot \|u\|_{L^\infty} \\
& \leq CMt \cdot \|\nabla u_t\|_{L^2} \cdot \|\nabla u\|_{H^1}^2.
\end{aligned}$$

It follows from Lemma 3.4 that

$$\begin{aligned}
(3.35) \quad & -2t \int \mu(\rho)_t \cdot d \cdot \nabla u_t dx \\
& \leq \frac{1}{8} \underline{\mu} \cdot t \|\nabla u_t\|_{L^2}^2 + \frac{CM^2}{\underline{\mu}} \cdot t (M_2^4 \bar{\rho}^2 \|\sqrt{\rho} u_t\|_{L^2}^4 + M_2^8 \bar{\rho}^8 \|\nabla u\|_{L^2}^{12}) \\
& \leq \frac{1}{8} \underline{\mu} \cdot t \|\nabla u_t\|_{L^2}^2 + \frac{CM^2 M_2^4 \bar{\rho}^2}{\underline{\mu}} \cdot t \|\sqrt{\rho} u_t\|_{L^2}^4 + \frac{CM^2 M_2^8 \bar{\rho}^8}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{12}.
\end{aligned}$$

Finally, taking into account the mass equation again, we arrive at

$$\begin{aligned}
(3.36) \quad & -t \int (\rho u)_t \cdot \nabla u \cdot u_t dx \\
& = -t \int \rho u \cdot \nabla (u \cdot \nabla u \cdot u_t) dx - t \int \rho u_t \cdot \nabla u \cdot u_t dx \\
& \leq t \int \rho |u| \cdot |\nabla u|^2 \cdot |u_t| dx + Ct \int \rho |u|^2 \cdot |\nabla^2 u| \cdot |u_t| dx \\
& \quad + t \int \rho |u|^2 \cdot |\nabla u| \cdot |\nabla u_t| dx + t \int \rho |u_t|^2 \cdot |\nabla u| dx \\
& = \sum_{i=1}^4 J_i.
\end{aligned}$$

Hence, it follows from Sobolev embedding inequality, Gagliardo-Nirenberg inequality, and Lemma 3.4 that

$$\begin{aligned}
 (3.37) \quad J_1 &\leq C\bar{\rho} \cdot t \|u_t\|_{L^6} \cdot \|u\|_{L^6} \cdot \|\nabla u\|_{L^3}^2 \\
 &\leq C\bar{\rho} \cdot t \|\nabla u_t\|_{L^2} \cdot \|\nabla u\|_{L^2}^2 \cdot \|\nabla u\|_{H^1} \\
 &\leq C\bar{\rho} \cdot t \|\nabla u_t\|_{L^2} \cdot \|\nabla u\|_{L^2}^2 \cdot \left(M_2 \bar{\rho}^{\frac{1}{2}} \|\sqrt{\rho} u_t\|_{L^2} + M_2^2 \bar{\rho}^2 \|\nabla u\|_{L^2}^3 \right) \\
 &\leq \frac{1}{8} \underline{\mu} \cdot t \|\nabla u_t\|_{L^2}^2 + \frac{CM_2^2 \bar{\rho}^3}{\underline{\mu}} \cdot t \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4 + \frac{CM_2^4 \bar{\rho}^6}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{10}.
 \end{aligned}$$

Similarly, it holds that

$$\begin{aligned}
 (3.38) \quad J_2 &\leq C\bar{\rho} \cdot t \|u_t\|_{L^6} \cdot \|\nabla^2 u\|_{L^2} \cdot \|u\|_{L^6}^2 \\
 &\leq C\bar{\rho} \cdot t \|\nabla u_t\|_{L^2} \cdot \|\nabla u\|_{H^1} \cdot \|\nabla u\|_{L^2}^2 \\
 &\leq \frac{1}{8} \underline{\mu} \cdot t \|\nabla u_t\|_{L^2}^2 + \frac{CM_2^2 \bar{\rho}^3}{\underline{\mu}} \cdot t \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4 + \frac{CM_2^4 \bar{\rho}^6}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{10},
 \end{aligned}$$

and

$$\begin{aligned}
 (3.39) \quad J_3 &\leq C\bar{\rho} \cdot t \|\nabla u_t\|_{L^2} \cdot \|\nabla u\|_{L^6} \cdot \|u\|_{L^6}^2 \\
 &\leq \frac{1}{8} \underline{\mu} \cdot t \|\nabla u_t\|_{L^2}^2 + \frac{CM_2^2 \bar{\rho}^3}{\underline{\mu}} \cdot t \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4 + \frac{CM_2^4 \bar{\rho}^6}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{10}.
 \end{aligned}$$

Owing to Lemma 3.4 and Sobolev embedding inequality,

$$\begin{aligned}
 (3.40) \quad J_4 &\leq Ct \|\sqrt{\rho} u_t\|_{L^4}^2 \cdot \|\nabla u\|_{L^2} \\
 &\leq Ct \|\sqrt{\rho} u_t\|_{L^2}^{\frac{1}{2}} \cdot \bar{\rho}^{\frac{3}{4}} \|\nabla u_t\|_{L^2}^{\frac{3}{2}} \cdot \|\nabla u\|_{L^2} \\
 &\leq \frac{1}{8} \underline{\mu} \cdot t \|\nabla u_t\|_{L^2}^2 + \frac{C\bar{\rho}^3}{\underline{\mu}} \cdot t \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4.
 \end{aligned}$$

Combine all the above estimates (3.33)-(3.40),

$$\begin{aligned}
 (3.41) \quad &\frac{d}{dt} t \int \rho |u_t|^2 dx + \underline{\mu} \cdot t \|\nabla u_t\|_{L^2}^2 \\
 &\leq \frac{C\bar{\rho}^3}{\underline{\mu}^3} \cdot t \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4 + \frac{CM_2^2 \bar{\rho}^3}{\underline{\mu}} \cdot t \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4 \\
 &\quad + \frac{CM_2^4 \bar{\rho}^6}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{10} + \frac{CM^2 M_2^4 \bar{\rho}^2}{\underline{\mu}} \cdot t \|\sqrt{\rho} u_t\|_{L^2}^4 \\
 &\quad + \frac{CM^2 M_2^8 \bar{\rho}^8}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{12} + \int \rho |u_t|^2 dx.
 \end{aligned}$$

Applying Gronwall's inequality,

$$\begin{aligned} & \sup_{t \in [0, T]} t \int \rho |u_t|^2 dx + \underline{\mu} \int_0^T t \|\nabla u_t\|_{L^2}^2 dt \\ & \leq \left[\int_0^T \left(\frac{CM_2^4 \bar{\rho}^6}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{10} + \frac{CM^2 M_2^8 \bar{\rho}^8}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{12} + \|\sqrt{\rho} u_t\|_{L^2}^2 \right) dt \right] \\ & \quad \cdot \exp \left\{ \int_0^T \left[\left(\frac{C \bar{\rho}^3}{\underline{\mu}^3} + \frac{CM_2^2 \bar{\rho}^3}{\underline{\mu}} \right) \|\nabla u\|_{L^2}^4 + \frac{CM^2 M_2^4 \bar{\rho}^2}{\underline{\mu}} \|\sqrt{\rho} u_t\|_{L^2}^2 \right] dt \right\}. \end{aligned}$$

Taking (3.20) and (3.23) into account,

$$\begin{aligned} & \sup_{t \in [0, T]} t \int \rho |u_t|^2 dx + \underline{\mu} \int_0^T t \|\nabla u_t\|_{L^2}^2 dt \\ (3.42) \quad & \leq \left[\int_0^T \left(\frac{CM_2^4 \bar{\rho}^6}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{10} + \frac{CM^2 M_2^8 \bar{\rho}^8}{\underline{\mu}} \cdot t \|\nabla u\|_{L^2}^{12} + \|\sqrt{\rho} u_t\|_{L^2}^2 \right) dt \right] \\ & \quad \cdot \exp \left\{ C \left(\frac{\bar{\rho}^4}{\underline{\mu}^4} + \frac{M_2^2 \bar{\rho}^4}{\underline{\mu}^2} + M^2 M_2^4 \bar{\rho}^2 \right) \right\}. \end{aligned}$$

According to Theorems 3.2, 3.5 and the assumption (3.18),

$$\begin{aligned} \int_0^T t \|\nabla u\|_{L^2}^{10} dt & \leq \sup_{t \in [0, T]} t \|\nabla u\|_{L^2}^2 \cdot \sup_{t \in [0, T]} \|\nabla u\|_{L^2}^6 \cdot \int_0^T \|\nabla u\|_{L^2}^2 dt \\ & \leq \frac{C \cdot \bar{\rho}^2}{\underline{\mu}^2} \|u_0\|_{L^2}^2 \leq \frac{C \cdot \bar{\rho}^2}{\underline{\mu}^2} \|\nabla u_0\|_{L^2}^2. \end{aligned}$$

Similarly,

$$\int_0^T t \|\nabla u\|_{L^2}^{12} dt \leq \frac{C \cdot \bar{\rho}^2}{\underline{\mu}^2} \|\nabla u_0\|_{L^2}^2.$$

And by virtue of Theorem 3.3,

$$\int_0^T \|\sqrt{\rho} u_t\|_{L^2}^2 dt \leq \underline{\mu} \cdot \|\nabla u_0\|_{L^2}^2.$$

Hence,

$$\begin{aligned} & \sup_{t \in [0, T]} t \int \rho |u_t|^2 dx + \underline{\mu} \int_0^T t \|\nabla u_t\|_{L^2}^2 dt \\ (3.43) \quad & \leq C \|\nabla u_0\|_{L^2}^2 \left(\frac{M_2^4 \bar{\rho}^8}{\underline{\mu}^3} + \frac{M^2 M_2^8 \bar{\rho}^{10}}{\underline{\mu}^3} + \underline{\mu} \right) \cdot \exp \left\{ C \left(\frac{\bar{\rho}^4}{\underline{\mu}^4} + \frac{M_2^2 \bar{\rho}^4}{\underline{\mu}^2} + M^2 M_2^4 \bar{\rho}^2 \right) \right\}. \end{aligned}$$

On the other hand, multiplying (3.41) by t , one has

$$\begin{aligned}
 & \frac{d}{dt} \left(\frac{t^2}{2} \int \rho |u_t|^2 dx \right) + \underline{\mu} t^2 \|\nabla u_t\|_{L^2}^2 \\
 & \leq \frac{C \bar{\rho}^3}{\underline{\mu}^3} \cdot t^2 \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4 + \frac{C M_2^2 \bar{\rho}^3}{\underline{\mu}} \cdot t^2 \|\sqrt{\rho} u_t\|_{L^2}^2 \cdot \|\nabla u\|_{L^2}^4 \\
 & \quad + \frac{C M_2^4 \bar{\rho}^6}{\underline{\mu}} \cdot t^2 \|\nabla u\|_{L^2}^{10} + \frac{C M^2 M_2^4 \bar{\rho}^2}{\underline{\mu}} \cdot t^2 \|\sqrt{\rho} u_t\|_{L^2}^4 \\
 & \quad + \frac{C M^2 M_2^8 \bar{\rho}^8}{\underline{\mu}} \cdot t^2 \|\nabla u\|_{L^2}^{12} + t \int \rho |u_t|^2 dx.
 \end{aligned} \tag{3.44}$$

Applying Gronwall's inequality,

$$\begin{aligned}
 & \sup_{t \in [0, T]} t^2 \int \rho |u_t|^2 dx + \underline{\mu} \int_0^T t^2 \|\nabla u_t\|_{L^2}^2 dt \\
 & \leq \left[\int_0^T \left(\frac{C M_2^4 \bar{\rho}^6}{\underline{\mu}} \cdot t^2 \|\nabla u\|_{L^2}^{10} + \frac{C M^2 M_2^8 \bar{\rho}^8}{\underline{\mu}} \cdot t^2 \|\nabla u\|_{L^2}^{12} + t \|\sqrt{\rho} u_t\|_{L^2}^2 \right) dt \right] \\
 & \quad \cdot \exp \left\{ C \left(\frac{\bar{\rho}^4}{\underline{\mu}^4} + \frac{M_2^2 \bar{\rho}^4}{\underline{\mu}^2} + M^2 M_2^4 \bar{\rho}^2 \right) \right\}.
 \end{aligned} \tag{3.45}$$

According to Theorems 3.2 and 3.5,

$$\begin{aligned}
 \int_0^T t^2 \|\nabla u\|_{L^2}^{10} dt & \leq \sup_{t \in [0, T]} t^2 \|\nabla u\|_{L^2}^4 \cdot \sup_{t \in [0, T]} \|\nabla u\|_{L^2}^4 \cdot \int_0^T \|\nabla u\|_{L^2}^2 dt \\
 & \leq \frac{C \bar{\rho}^3}{\underline{\mu}^3} \|\nabla u_0\|_{L^2}^2.
 \end{aligned}$$

Similarly,

$$\int_0^T t^2 \|\nabla u\|_{L^2}^{12} dt \leq \frac{C \bar{\rho}^3}{\underline{\mu}^3} \|\nabla u_0\|_{L^2}^2.$$

Hence,

$$\begin{aligned}
 & \sup_{t \in [0, T]} t^2 \int \rho |u_t|^2 dx + \underline{\mu} \int_0^T t^2 \|\nabla u_t\|_{L^2}^2 dt \\
 & \leq C \|\nabla u_0\|_{L^2}^2 \left(\frac{M_2^4 \bar{\rho}^9}{\underline{\mu}^4} + \frac{M^2 M_2^8 \bar{\rho}^{11}}{\underline{\mu}^4} + \bar{\rho} \right) \cdot \exp \left\{ C \left(\frac{\bar{\rho}^4}{\underline{\mu}^4} + \frac{M_2^2 \bar{\rho}^4}{\underline{\mu}^2} + M^2 M_2^4 \bar{\rho}^2 \right) \right\}.
 \end{aligned}$$

□

Lemma 3.7. *Suppose (ρ, u, P) is the unique local strong solution to (1.1) on $[0, T]$, with the initial data (ρ_0, u_0) , and satisfies the assumptions (3.17)-(3.19).*

Then for any $r \in (3, \min\{q, 6\})$

$$(3.47) \quad \begin{aligned} & \int_0^T \|\nabla u\|_{L^\infty} dt \\ & \leq C \|\nabla u_0\|_{L^2} \left[M_r \bar{\rho}^{\frac{5r-6}{4r}} \underline{\mu}^{-\frac{3(r-2)}{4r}} \left(1 + \frac{\bar{\rho}}{\underline{\mu}}\right)^{\frac{1}{2}} \Theta_1^{\frac{1}{2}} \exp\{C\Theta_2\} + M_r^{\frac{5r-6}{r}} \frac{\bar{\rho}^{\frac{6(r-1)}{r}}}{\underline{\mu}} \right]. \end{aligned}$$

Proof. By virtue of Lemma 2.1, one has for $r \in (3, \min\{q, 6\})$

$$(3.48) \quad \begin{aligned} \|\nabla u\|_{W^{1,r}} & \leq CM_r (\|\rho u_t\|_{L^r} + \|\rho u \cdot \nabla u\|_{L^r}) \\ & \leq CM_r \left(\|\rho u_t\|_{L^2}^{\frac{6-r}{2r}} \cdot \|\rho u_t\|_{L^6}^{\frac{3(r-2)}{2r}} + \bar{\rho} \|u\|_{L^6} \cdot \|\nabla u\|_{L^{6r/(6-r)}} \right) \\ & \leq CM_r \left(\|\rho u_t\|_{L^2}^{\frac{6-r}{2r}} \cdot \|\rho u_t\|_{L^6}^{\frac{3(r-2)}{2r}} + \bar{\rho} \|\nabla u\|_{L^2}^{\frac{6(r-1)}{5r-6}} \cdot \|\nabla u\|_{W^{1,r}}^{\frac{4r-6}{5r-6}} \right). \end{aligned}$$

Applying Young's inequality and Sobolev embedding inequality,

$$(3.49) \quad \|\nabla u\|_{W^{1,r}} \leq CM_r \bar{\rho}^{\frac{5r-6}{4r}} \cdot \|\sqrt{\rho} u_t\|_{L^2}^{\frac{6-r}{2r}} \cdot \|\nabla u_t\|_{L^2}^{\frac{3(r-2)}{2r}} + CM_r^{\frac{5r-6}{r}} \bar{\rho}^{\frac{5r-6}{r}} \cdot \|\nabla u\|_{L^2}^{\frac{6(r-1)}{r}}.$$

Hence,

$$(3.50) \quad \begin{aligned} & \int_0^T \|\nabla u\|_{L^\infty} dt \\ & \leq C \int_0^T \|\nabla u\|_{W^{1,r}} dt \\ & \leq C \int_0^T \left(M_r \bar{\rho}^{\frac{5r-6}{4r}} \|\sqrt{\rho} u_t\|_{L^2}^{\frac{6-r}{2r}} \|\nabla u_t\|_{L^2}^{\frac{3(r-2)}{2r}} + M_r^{\frac{5r-6}{r}} \bar{\rho}^{\frac{5r-6}{r}} \|\nabla u\|_{L^2}^{\frac{6(r-1)}{r}} \right) dt. \end{aligned}$$

If $T \leq 1$, according to Theorem 3.6,

$$\begin{aligned} & \int_0^T \|\sqrt{\rho} u_t\|_{L^2}^{\frac{6-r}{2r}} \cdot \|\nabla u_t\|_{L^2}^{\frac{3(r-2)}{2r}} dt \\ & \leq \int_0^T \left(t^{\frac{1}{2}} \|\sqrt{\rho} u_t\|_{L^2} \right)^{\frac{6-r}{2r}} \cdot \left(t^{\frac{1}{2}} \|\nabla u_t\|_{L^2} \right)^{\frac{3(r-2)}{2r}} \cdot t^{-\frac{1}{2}} dt \\ & \leq C \left(\sup_{t \in [0, T]} t^{\frac{1}{2}} \|\sqrt{\rho} u_t\|_{L^2} \right)^{\frac{6-r}{2r}} \cdot \left(\int_0^T t \|\nabla u_t\|_{L^2}^2 dt \right)^{\frac{3(r-2)}{4r}} \left(\int_0^T t^{-\frac{2r}{r+6}} dt \right)^{\frac{r+6}{4r}} \\ & \leq C \underline{\mu}^{-\frac{3(r-2)}{4r}} \cdot \|\nabla u_0\|_{L^2}^2 \Theta_1^{\frac{1}{2}} \exp\{C\Theta_2\}. \end{aligned}$$

If $T > 1$, applying Theorem 3.6 again,

$$\begin{aligned}
& \int_0^T \|\sqrt{\rho}u_t\|_{L^2}^{\frac{6-r}{2r}} \cdot \|\nabla u_t\|_{L^2}^{\frac{3(r-2)}{2r}} dt \\
& \leq \int_0^T \left(t^{\frac{1}{2}}\|\sqrt{\rho}u_t\|_{L^2}\right)^{\frac{6-r}{2r}} \cdot \left(t^{\frac{1}{2}}\|\nabla u_t\|_{L^2}\right)^{\frac{3(r-2)}{2r}} \cdot t^{-\frac{1}{2}} dt \\
& \quad + \int_1^T (t\|\sqrt{\rho}u_t\|_{L^2})^{\frac{6-r}{2r}} \cdot (t\|\nabla u_t\|_{L^2})^{\frac{3(r-2)}{2r}} \cdot t^{-1} dt \\
& \leq C \left(\sup_{t \in [0, T]} t^{\frac{1}{2}}\|\sqrt{\rho}u_t\|_{L^2}\right)^{\frac{6-r}{2r}} \cdot \left(\int_0^T t\|\nabla u_t\|_{L^2}^2 dt\right)^{\frac{3(r-2)}{4r}} \left(\int_0^T t^{-\frac{2r}{r+6}} dt\right)^{\frac{r+6}{4r}} \\
& \quad + C \left(\sup_{t \in [0, T]} t\|\sqrt{\rho}u_t\|_{L^2}\right)^{\frac{6-r}{2r}} \cdot \left(\int_0^T t^2\|\nabla u_t\|_{L^2}^2 dt\right)^{\frac{3(r-2)}{4r}} \left(\int_1^T t^{-\frac{4r}{r+6}} dt\right)^{\frac{6+r}{4r}} \\
& \leq C \underline{\mu}^{-\frac{3(r-2)}{4r}} \|\nabla u_0\|_{L^2} \Theta_1^{\frac{1}{2}} \exp\{C\Theta_2\} + C \underline{\mu}^{-\frac{3(r-2)}{4r}} \left(\frac{\bar{\rho}}{\underline{\mu}}\right)^{\frac{1}{2}} \|\nabla u_0\|_{L^2} \Theta_1^{\frac{1}{2}} \exp\{C\Theta_2\} \\
& \leq C \underline{\mu}^{-\frac{3(r-2)}{4r}} \left(1 + \frac{\bar{\rho}}{\underline{\mu}}\right)^{\frac{1}{2}} \|\nabla u_0\|_{L^2} \Theta_1^{\frac{1}{2}} \exp\{C\Theta_2\}.
\end{aligned}$$

On the other hand,

$$(3.51) \quad \int_0^T \|\nabla u\|_{L^2}^{\frac{6(r-1)}{r}} dt \leq \sup_{t \in [0, T]} \|\nabla u\|_{L^2}^{\frac{4r-6}{r}} \cdot \int_0^T \|\nabla u\|_{L^2}^2 dt \leq \frac{C \cdot \bar{\rho}}{\underline{\mu}} \|\nabla u_0\|_{L^2}.$$

Therefore,

$$\begin{aligned}
(3.52) \quad & \int_0^T \|\nabla u\|_{L^\infty} dt \\
& \leq C \|\nabla u_0\|_{L^2} \left[M_r \bar{\rho}^{\frac{5r-6}{4r}} \underline{\mu}^{-\frac{3(r-2)}{4r}} \left(1 + \frac{\bar{\rho}}{\underline{\mu}}\right)^{\frac{1}{2}} \Theta_1^{\frac{1}{2}} \exp\{C\Theta_2\} + M_r^{\frac{5r-6}{r}} \frac{\bar{\rho}^{\frac{6(r-1)}{r}}}{\underline{\mu}} \right] \\
& \triangleq C_2(M, \bar{\rho}, \underline{\mu}, \bar{\mu}) \|\nabla u_0\|_{L^2}.
\end{aligned}$$

□

Theorem 3.8. *Suppose (ρ, u, P) is the unique local strong solution to (1.1) on $[0, T]$ and the assumptions (3.17)-(3.19). There exists a positive number ϵ_0 , depending on $\Omega, q, M, \bar{\rho}, \bar{\mu}, \underline{\mu}$ such that if*

$$\|\nabla u_0\|_{L^2} \leq \epsilon_0,$$

then

$$(3.53) \quad \sup_{t \in [0, T]} \|\nabla \mu(\rho)\|_{L^q} \leq 2M,$$

and

$$\sup_{t \in [0, T]} \|\nabla \rho\|_{L^q} \leq 2\|\nabla \rho_0\|_{L^q}.$$

Proof. Consider the x_i -derivative of the renormalized mass equation for $\mu(\rho)$,

$$(\partial_i \mu(\rho))_t + (\partial_i u \cdot \nabla) \mu(\rho) + u \cdot \nabla \partial_i \mu(\rho) = 0.$$

It implies that for every $t \in [0, T]$, in view of Lemma 3.7, one has

$$\begin{aligned} (3.54) \quad \|\nabla \mu(\rho)(t)\|_{L^q} &\leq \|\nabla \mu(\rho_0)\|_{L^q} \cdot \exp \left\{ \int_0^t \|\nabla u\|_{L^\infty} ds \right\} \\ &\leq \|\nabla \mu(\rho_0)\|_{L^q} \cdot \exp \left\{ C_2(M, \bar{\rho}, \underline{\mu}, \bar{\mu}) \cdot \|\nabla u_0\|_{L^2} \right\}. \end{aligned}$$

Choose some small positive constant ϵ_0 , which satisfies

$$\epsilon_0 \leq \frac{1}{2}, \quad \epsilon_0^2 \cdot C_1 \underline{\mu}^{-2} (M_2^2 + M^4 M_2^6) \bar{\rho}^4 \leq \ln 2,$$

and

$$\epsilon_0 \cdot C_2(M, \bar{\rho}, \bar{\mu}, \underline{\mu}) \leq \ln 2.$$

If $\|\nabla u_0\|_{L^2} \leq \epsilon_0$, (3.53) holds.

Similarly,

$$\begin{aligned} (3.55) \quad \|\nabla \rho(t)\|_{L^q} &\leq \|\nabla \rho_0\|_{L^q} \cdot \exp \left\{ \int_0^t \|\nabla u\|_{L^\infty} ds \right\} \\ &\leq 2 \|\nabla \rho_0\|_{L^q}, \end{aligned}$$

Therefore, Theorem 3.8 is proved. $\square$

3.2. Proof of Theorem 1.2. With the a priori estimates in Subsection 3.1 in hand, we are prepared for the proof of Theorem 1.2.

Proof. According to Theorem 1.1, there exists a $T_* > 0$ such that the density-dependent Navier-Stokes system (1.1) has a unique local strong solution (ρ, u, P) on $[0, T_*]$. We plan to extend the local solution to a global one.

Since $\|\nabla \mu(\rho_0)\|_{L^q} = M < 4M$, and due to the continuity of $\nabla \mu(\rho)$ in L^q and ∇u_0 in L^2 , there exists a $T_1 \in (0, T_*)$ such that $\sup_{0 \leq t \leq T_1} \|\nabla \mu(\rho)(t)\|_{L^q} \leq 4M$, and at the same time $\sup_{0 \leq t \leq T_1} \|\nabla u(t)\|_{L^2} \leq 2\|\nabla u_0\|_{L^2}$.

Set

$$\begin{aligned} T^* &= \sup \{T \mid (\rho, u, P) \text{ is a strong solution to (1.1) on } [0, T]\} \\ T_1^* &= \sup \left\{ T \mid \begin{array}{l} (\rho, u, P) \text{ is a strong solution to (1.1) on } [0, T], \\ \sup_{0 \leq t \leq T} \|\nabla \mu(\rho)\|_{L^q} \leq 4M, \text{ and } \sup_{0 \leq t \leq T} \|\nabla u\|_{L^2} \leq 2\|\nabla u_0\|_{L^2}. \end{array} \right\}. \end{aligned}$$

Then $T_1^* \geq T_1 > 0$. Recalling Theorems 3.3 and 3.8, it's easy to verify

$$(3.56) \quad T^* = T_1^*,$$

provided that $\|\nabla u_0\|_{L^2} \leq \epsilon_0$ as assumed.

We claim that $T^* = \infty$. Otherwise, assuming that $T^* < \infty$. By virtue of Theorems 3.3 and 3.8, for every $t \in [0, T^*)$, it holds that

$$(3.57) \quad \|\nabla \rho(t)\|_{L^q} \leq 2\|\nabla \rho_0\|_{L^q}, \quad \text{and} \quad \|\nabla u(t)\|_{L^2} \leq \sqrt{2}\|\nabla u_0\|_{L^2}.$$

which contradicts to the blowup criterion (1.5). Hence we complete the proof for Theorem 1.2. $\square$

Acknowledgement: The research of Xiangdi Huang is supported in part by NSFC Grant No. 11101392. The research of Yun Wang is supported in part by NSFC Grant No. 11301365.

REFERENCES

- [1] H. Abidi, G. L. Gui and P. Zhang, On the decay and stability to global solutions of the 3-D inhomogeneous Navier-Stokes equations, *Comm. Pure Appl. Math.*, **64** (2011), 832-881.
- [2] H. Abidi, G. Gui and P. Zhang, On the wellposedness of three dimensional inhomogeneous Navier-Stokes equations in the critical spaces, *Arch. Ration. Mech. Anal.*, **204** (2012), 189-230.
- [3] H. Abidi, P. Zhang, On the global well-posedness of 2-D density-dependent Navier-Stokes system with variable viscosity. <http://arxiv.org/abs/1301.2371>.
- [4] S. A. Antontsev and A. V. Kazhikov, Mathematical Study of Flows of Nonhomogeneous Fluids; Lecture Notes, Novosibirsk State University, Novosibirsk, U.S.S.R., 1973(Russian).
- [5] S. A. Antontsev, A. V. Kazhikov and V. N. Monakhov, Boundary Value Problems in Mechanics of Nonhomogeneous Fluids, *North-Holland, Amsterdam*, 1990.
- [6] Y. Cho and H. Kim, Unique solvability for the density-dependent Navier-Stokes equations, *Nonlinear Anal.* **59** (2004), no. 4, 465-489.
- [7] H. Y. Choe and H. Kim, Strong solutions of the NavierStokes equations for nonhomogeneous incompressible fluids, *Comm. Partial Differential Equations*, **28** (2003) no. 5-6, 1183-1201.
- [8] W. Craig, H. X. Huang and Y. Wang, Global strong solutions fo 3D nonhomogeneous incompressible Navier-Stokes equations, *J. Math. Fluid Mech.*, **15** (2013), 747-758.
- [9] R. Danchin, Density-dependent incompressible viscous fluids in critical spaces, *Proc. Roy. Soc. Edinburgh Sect. A*, **133** (2003), 1311-1334.
- [10] R. Danchin, Local and global well-posedness results for flows of inhomogeneous viscous fluids, *Adv. Differential Equations*, **9** (2004), 353-386.
- [11] R. Danchin and P. B. Mucha, A Lagrangian approach for the incompressible Navier-Stokes equations with variable density, *Comm. Pure. Appl. Math.*, **65** (2012), 1458-1480.
- [12] R. Danchin and P. B. Mucha, Incompressible flows with piecewise constant density. *Arch. Ration. Mech. Anal.* **207** (2013), no. 3, 991-1023.
- [13] B. Desjardins, Regularity results for two-dimensional flows of multiphase viscous fluids, *Arch. Rat. Mech. Anal.*, **137** (1997), 135-158.
- [14] R. J. DiPerna and P.L. Lions, Equations différentielles ordinaires et équations de transport avec des coefficients irréguliers, in: Séminaire EDP 1988-1989, Ecole Polytechnique, Palaiseau, 1989.
- [15] G. P. Galdi, An Introduction to the Mathematical Theory of the Navier-Stokes Equations, vol. I, revised edition, Springer Tracts in Natural Philosophy (ed. C. Truesdell), **38**, Springer-Verlag, Berlin, 1998.
- [16] M. Giaquinta, G. Modica, Nonlinear systems of the type of the stationary Navier-Stokes system, *J. Reine Angew. Math.*, **330** (1982), 173-214.
- [17] D. Hoff, Global solutions of the Navier-Stokes equations for multidimensional compressible flow with discontinuous initial data, *J. Differential Equations*, **120** (1995), no. 1, 215-254.
- [18] J. C. Huang, M. Paicu and P. Zhang, Global solutions to 2-D inhomogeneous Navier-Stokes system with general velocity, *J. Math. Pures Appl. (9)*, **100** (2013), no. 6, 806-831.
- [19] X. D. Huang, J. Li and Z. P. Xin, Global well-posedness of classical solutions with large oscillations and vacuum to the three-dimensional isentropic compressible Navier-Stokes equations, *Comm. Pure Appl. Math.*, **65** (2012), no. 4, 549-585.
- [20] X. D. Huang and Y. Wang, Global Strong Solution to the 2D Nonhomogeneous Incompressible MHD System, *J. Differential Equations*, **254** (2013), 511-527.
- [21] A. V. Kazhikov, Resolution of boundary value problems for nonhomogeneous viscous fluids, *Dokl. Akad. Nauk.*, **216** (1974), 1008-1010.

- [22] O. Ladyzhenskaya and V. A. Solonnikov, Unique solvability of an initial and boundary value problem for viscous incompressible non-homogeneous fluids, *J. Soviet Math.*, **9** (1978), 697-749.
- [23] P. L. Lions, Mathematical Topics in Fluid Mechanics, Vol. I: Incompressible Models, *Oxford Lecture Series in Math. and Its Appl.* **3**, Oxford University Press, New York, 1996.
- [24] M. Paicu and P. Zhang, Global solutions to the 3-D incompressible inhomogeneous Navier-Stokes system, *J. Functional Anal.*, **262** (2012), 3556-3584.
- [25] R. Salvi, The equations of viscous incompressible non-homogeneous fluids: on the existence and regularity, *J. Austral. Math. Soc. Ser. B*, **33** (1991), 94-110.
- [26] J. Simon, Nonhomogeneous viscous incompressible fluids: Existence of velocity, density, and pressure, *SIAM J. Math. Anal.*, **21** (1990), 1093-1117.

NCMIS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CAS, BEIJING 100190,
P. R. CHINA

E-mail address: `xdhuang@amss.ac.cn`

DEPARTMENT OF MATHEMATICS, SOOCHOW UNIVERSITY, 1 SHIZI STREET, SUZHOU 215006,
P.R. CHINA

E-mail address: `ywang3@suda.edu.cn`